

WAIPUKURAU FACIAL ECZEMA SPORE COUNT MONITOR - 2019

Please note that the following spore counts are individual counts from selected properties and are only a rough guide as to counts in the area. Individual paddocks or geographical areas can vary hugely depending on local topography/climate and pasture type/condition. For an accurate assessment of your pastures regular spore counting is recommended during danger periods.

As a guide levels of over 30,000 spores/gram of pasture will cause toxicity if grazed for over 2 weeks while levels of over 100,000 spores/gram can cause serious liver pathology after as little as 3 days.

LOCATION	February				March				April				May				
	Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4	Week 5
Hinerangi	15,000	5,000	30,000	15,000	30,000	5,000	200,000	110,000	25,000	75,000	20,000	25,000	30,000	0			
Patangata		5,000	5,000	25,000		40,000	40,000	45,000									
Ashley Clinton	15,000	20,000	5,000	25,000	40,000	30,000	45,000	50,000	20,000								
Porangahau Rd	0	5,000	0	5,000	5,000	0	20,000	10,000	30,000	150,000	5,000	0					
Mangatarata	10,000 20,000	10,000	35,000 5,000	65,000 45,000	30,000 55,000	20,000 65,000	50,000 95,000	20,000 35,000	5,000 10,000		10,000 15,000	5,000 25,000	10,000 25,000				
Wilder Rd	10,000	5,000	0	5,000	5,000	30,000	15,000	20,000	10,000	0	5,000	15,000	5,000	5,000			
Motere Rd						5,000	40,000 10,000	40,000 0	10,000 65,000	5,000							
Takapau								0									
Elsthorpe	10,000	20,000	15,000		50,000	55,000	55,000	35,000	35,000	20,000	20,000						
Wallingford	5,000	5,000	5,000	15,000	5,000	65,000	260,000	35,000	35,000	25,000	45,000	70,000	25,000				
Omakere	5,000	0	35,000	10,000	15,000	45,000	95,000	25,000	5,000	5,000	35,000		15,000	20,000			
Farm Rd			10,000	20,000 25,000	20,000 25,000	25,000 50,000	65,000 85,000	15,000 40,000	15,000 15,000								
Onga Onga								30,000 150,000	10,000 15,000	95,000 130,000	25,000 35,000	30,000 45,000	10,000 30,000	5,000 10,000			