VETservices Hawke's Bay

WAIPUKURAU FACIAL ECZEMA SPORE COUNT MONITOR 2020

Please note that the following spore counts are individual counts from selected properties and are only a rough guide as to counts in the area. Individual paddocks or geographical areas can vary hugely depending on local topography/climate and pasture type/condition. For an accurate assessment of your pastures regular spore counting is recommended during danger periods.

As a guide levels of over 30,000 spores/gram of pasture will cause toxicity if grazed for over 2 weeks while levels of over 100,000 spores/gram can cause serious liver pathology after as little as 3 days.

LOCATION	March				April				May				June					July
	Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4	Week 5	Week 1
Hinerangi	35,000	30,000	20,000	20,000														
Patangata																		
Ashley Clinton	300,000	175,000	65,000	205,000	585,000													
Porangahau	10,000																	
Mangatarata	10,000	0	5,000															
Wilder Rd	0																	
Motere Rd																		
Takapau																		
Elsthorpe		20,000		5,000	35,000													
Wallingford	10,000	5,000	5,000															
Omakere	105,000	0	5,000															
Farm Rd		5,000																
Lindsay Rd	0	0	0		0													