

### HASTINGS FACIAL ECZEMA SPORE COUNT MONITOR - 2016

Please note that the following spore counts are individual counts from selected properties and are only a rough guide as to counts in the area. Individual paddocks or geographical areas can vary hugely depending on local topography/climate and pasture type/condition. For an accurate assessment of your pastures regular spore counting is recommended during danger periods.

As a guide levels of over 30,000 spores/gram of pasture will cause toxicity if grazed for over 2 weeks while levels of over 100,000 spores/gram can cause serious liver pathology after as little as 3 days

LOCATION	January				February				Week 1	March			
	Week 1	Week 2	Week 3	Week 4	Week 1	Week 2	Week 3	Week 4		Week 2	Week 3	Week 4	Week 5
TUTIRA				0	15,000	5,000	5,000						
KERERU													
PATOKA			25,000 10,000		15,000 20,000	65,000	10,000 75,000	20,000 65,000	15,000 45,000	20,000 40,000	30,000	30,000	40,000
BAYVIEW			10,000	20,000	55,000 50,000	20,000 25,000	30,000 30,000		50,000 75,000	115,000 90,000	35,000 140,000	15,000 140,000	105,000 325,000
DARTMOOR RD				30,000		15,000							
WAIMARAMA													
TAIHAPE (whana Whana)					5,000 15,000	5,000 15,000	10,000 20,000	5,000 5,000		50,000	40,000		15,000 75,000
POUKAWA						5,000							
NGATARAWA					20,000 70,000 15,000	15,000 15,000	20,000						0 25,000
TE POHUE							5,000 5,000						

